

Nickel at Lockwood

This page was updated on October 31, 2003

Summary Statistics				
<i>Month</i>	<i>Number of Samples</i>	<i>Minimum (ng/m3)</i>	<i>Maximum (ng/m3)</i>	<i>Average (ng/m3)</i>
November-01	1	5.7	5.7	5.7
December-01	3	2.6	8.1	4.4
January-02	3	1.0	2.9	2.0
February-02	5	1.0	6.2	3.4
March-02	5	1.0	7.0	3.7
April-02	5	1.0	3.2	2.1
May-02	4	2.3	4.7	3.4
June-02	5	2.2	5.9	3.8
July-02	5	3.0	6.0	4.2
August-02	-	no data	no data	no data
September-02	-	no data	no data	no data
October-02	-	no data	no data	no data
November-02	4	6.0	9.0	6.8
December-02	4	0.5	10.0	3.6
January-03	4	3.0	5.0	4.0
February-03	5	2.0	4.0	3.2
March-03	-	no data	no data	no data
April-03	-	no data	no data	no data
SUMMARY	Total	Minimum	Maximum	Average
	53	0.5	10.0	3.9

* The lowest concentration that can be measured (LOD) is 2 nanograms per cubic meter (ng/m3) through June 2002, and starting in July 2002 the LOD is 1 ng/m3. Values below the LOD are set to 0.5 * LOD.